

Problem 7

In each of Problems 1 through 12:

- Draw a direction field for the given differential equation.
- Based on an inspection of the direction field, describe how solutions behave for large t .
- Find the general solution of the given differential equation, and use it to determine how solutions behave as $t \rightarrow \infty$.

$$y' + 2ty = 2te^{-t^2}$$

Solution

The direction field is a two-dimensional vector field that shows what the direction of the solution is at every point in a region. Every solution to the differential equation is a curve drawn such that the direction field vectors are tangent to it at every point.

$$\langle dt, dy \rangle = \left\langle 1, \frac{dy}{dt} \right\rangle dt = \langle 1, -2ty + 2te^{-t^2} \rangle dt$$




Figure 1: In blue are the direction field vectors and in red are possible solutions to the differential equation, depending what the initial condition is. All solutions appear to converge to $y = 0$ as $t \rightarrow \infty$.

$$y' + 2ty = 2te^{-t^2}$$

This is a first-order linear inhomogeneous equation, so it can be solved with an integrating factor.

$$I(t) = \exp\left(\int^t 2s \, ds\right) = e^{t^2}$$

Multiply both sides of the ODE by $I(t)$.

$$e^{t^2} y' + 2te^{t^2} y = 2t$$

The left side can be written as $d/dt(Iy)$ by the product rule.

$$\frac{d}{dt}(e^{t^2} y) = 2t$$

Integrate both sides with respect to t .

$$e^{t^2} y = t^2 + C$$

Divide both sides by e^{t^2} to solve for $y(t)$.

$$y(t) = e^{-t^2}(t^2 + C)$$

$y(t)$ vanishes as $t \rightarrow \infty$.

$$\lim_{t \rightarrow \infty} y(t) = 0$$