

Problem 10

In each of Problems 9 through 20:

- Find the solution of the given initial value problem in explicit form.
- Plot the graph of the solution.
- Determine (at least approximately) the interval in which the solution is defined.

$$y' = (1 - 2x)/y, \quad y(1) = -2$$

Solution

Part (a)

This ODE is separable because it is of the form $y' = f(x)g(y)$, so it can be solved by separating variables.

$$\frac{dy}{dx} = \frac{1 - 2x}{y}$$

Bring the terms with y to the left and bring the terms with x to the right.

$$y \, dy = (1 - 2x) \, dx$$

Integrate both sides.

$$\int y \, dy = \int (1 - 2x) \, dx$$
$$\frac{y^2}{2} = x - x^2 + C$$

Now apply the initial condition $y(1) = -2$ to determine C .

$$\frac{(-2)^2}{2} = 1 - 1^2 + C \quad \rightarrow \quad 2 = C$$

The previous equation then becomes


$$\frac{y^2}{2} = x - x^2 + 2$$
$$y^2 = -2(x^2 - x - 2)$$
$$y(x) = \pm \sqrt{-2(x - 2)(x + 1)}.$$

We choose the minus sign so that the initial condition $y(1) = -2$ is satisfied. Therefore, the solution to the initial value problem is

$$y(x) = -\sqrt{2(2 - x)(x + 1)}.$$

Part (b)

Below is a plot of $y(x)$ versus x .

**Part (c)**

The solution we found is only valid along the curve that passes through $x = 1$ because $x = 1$ is where the initial condition is given. It is valid over $-1 < x < 2$, the domain of y .