

Problem 18

In each of Problems 17 through 22, find the solution of the given initial value problem. Sketch the graph of the solution and describe its behavior for increasing t .

$$y'' + 4y' + 5y = 0, \quad y(0) = 1, \quad y'(0) = 0$$

Solution

Since this is a linear homogeneous constant-coefficient ODE, the solution is of the form $y = e^{rt}$.

$$y = e^{rt} \quad \rightarrow \quad y' = re^{rt} \quad \rightarrow \quad y'' = r^2e^{rt}$$

Substitute these expressions into the ODE.

$$r^2e^{rt} + 4(re^{rt}) + 5(e^{rt}) = 0$$

Divide both sides by e^{rt} .

$$\begin{aligned} r^2 + 4r + 5 &= 0 \\ r &= \frac{-4 \pm \sqrt{16 - 4(5)(1)}}{2} = \frac{-4 \pm \sqrt{-4}}{2} = \frac{-4 \pm 2i}{2} = -2 \pm i \\ r &= \{-2 - i, -2 + i\} \end{aligned}$$

Two solutions to the ODE are $y = e^{(-2-i)t}$ and $y = e^{(-2+i)t}$, so the general solution is a linear combination of the two.

$$\begin{aligned} y(t) &= C_1e^{(-2-i)t} + C_2e^{(-2+i)t} \\ &= C_1e^{-2t-it} + C_2e^{-2t+it} \\ &= C_1e^{-2t}e^{-it} + C_2e^{-2t}e^{it} \\ &= C_1e^{-2t}[\cos(-t) + i\sin(-t)] + C_2e^{-2t}[\cos(t) + i\sin(t)] \\ &= C_1e^{-2t}(\cos t - i\sin t) + C_2e^{-2t}(\cos t + i\sin t) \\ &= C_1e^{-2t}\cos t - iC_1e^{-2t}\sin t + C_2e^{-2t}\cos t + iC_2e^{-2t}\sin t \\ &= (C_1 + C_2)e^{-2t}\cos t + (-iC_1 + iC_2)e^{-2t}\sin t \end{aligned}$$

Using C_3 for $C_1 + C_2$ and C_4 for $-iC_1 + iC_2$, the real general solution is

$$y(t) = C_3e^{-2t}\cos t + C_4e^{-2t}\sin t.$$

Take a derivative of it.


$$y'(t) = -2C_3e^{-2t}\cos t - C_3e^{-2t}\sin t - 2C_4e^{-2t}\sin t + C_4e^{-2t}\cos t$$

Apply the initial conditions now to determine C_3 and C_4 .

$$\begin{aligned} y(0) &= C_3 = 1 \\ y'(0) &= -2C_3 + C_4 = 0 \end{aligned}$$

This system of equations yields $C_3 = 1$ and $C_4 = 2$. Therefore,

$$y(t) = e^{-2t}\cos t + 2e^{-2t}\sin t.$$



The solution exponentially decays to zero.