

Exercise 5

Use the function

$$f(z) = \frac{z^{1/3}}{(z+a)(z+b)} = \frac{e^{(1/3)\log z}}{(z+a)(z+b)} \quad (|z| > 0, 0 < \arg z < 2\pi)$$

and a closed contour similar to the one in Fig. 103 (Sec. 84) to show formally that

$$\int_0^\infty \frac{\sqrt[3]{x}}{(x+a)(x+b)} dx = \frac{2\pi}{\sqrt{3}} \cdot \frac{\sqrt[3]{a} - \sqrt[3]{b}}{a-b} \quad (a > b > 0).$$

Solution

In order to evaluate this integral, consider the given function in the complex plane and the contour in Fig. 103. Singularities occur where the denominator is equal to zero.

$$\begin{aligned} (z+a)(z+b) &= 0 \\ z &= -a \quad \text{or} \quad z = -b \end{aligned}$$

Since $z^{1/3}$ can be written in terms of $\log z$, a branch cut for the function needs to be chosen.

$$z^{1/3} = \exp\left(\frac{1}{3}\log z\right)$$

It has been chosen here to be the axis of positive real numbers.

$$\begin{aligned} &= \exp\left[\frac{1}{3}(\ln r + i\theta)\right], \quad (|z| > 0, 0 < \theta < 2\pi) \\ &= r^{1/3} e^{i\theta/3}, \end{aligned}$$

where $r = |z|$ is the magnitude of z and $\theta = \arg z$ is the argument of z .




Figure 1: This is essentially Fig. 103 with the singularities at $z = -a$ and $z = -b$ marked. The squiggly line represents the branch cut ($|z| > 0, 0 < \theta < 2\pi$).

According to Cauchy's residue theorem, the integral of $z^{1/3}/[(z+a)(z+b)]$ around the closed contour is equal to $2\pi i$ times the sum of the residues at the enclosed singularities.

$$\oint_C \frac{z^{1/3}}{(z+a)(z+b)} dz = 2\pi i \left[\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} + \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} \right]$$

This closed loop integral is the sum of four integrals, one over each arc in the loop.

$$\begin{aligned} \int_{L_1} \frac{z^{1/3}}{(z+a)(z+b)} dz + \int_{L_2} \frac{z^{1/3}}{(z+a)(z+b)} dz + \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz + \int_{C_R} \frac{z^{1/3}}{(z+a)(z+b)} dz \\ = 2\pi i \left[\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} + \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} \right] \quad (1) \end{aligned}$$

The parameterizations for the arcs are as follows.

$$\begin{aligned} L_1: \quad z &= re^{i0}, & r &= \rho \quad \rightarrow \quad r = R \\ L_2: \quad z &= re^{i2\pi}, & r &= R \quad \rightarrow \quad r = \rho \\ C_\rho: \quad z &= \rho e^{i\theta}, & \theta &= 2\pi \quad \rightarrow \quad \theta = 0 \\ C_R: \quad z &= R e^{i\theta}, & \theta &= 0 \quad \rightarrow \quad \theta = 2\pi \end{aligned}$$

As a result,

$$\begin{aligned} \int_{L_1} \frac{z^{1/3}}{(z+a)(z+b)} dz + \int_{L_2} \frac{z^{1/3}}{(z+a)(z+b)} dz &= \int_\rho^R \frac{(re^{i0})^{1/3}}{(re^{i0}+a)(re^{i0}+b)} (dr e^{i0}) + \int_R^\rho \frac{(re^{i2\pi})^{1/3}}{(re^{i2\pi}+a)(re^{i2\pi}+b)} (dr e^{i2\pi}) \\ &= \int_\rho^R \frac{r^{1/3}}{(r+a)(r+b)} dr + \int_R^\rho \frac{r^{1/3} e^{i2\pi/3}}{(r+a)(r+b)} dr \\ &= \int_\rho^R \frac{r^{1/3}}{(r+a)(r+b)} dr - \int_\rho^R \frac{r^{1/3} e^{i2\pi/3}}{(r+a)(r+b)} dr \\ &= (1 - e^{2i\pi/3}) \int_\rho^R \frac{r^{1/3}}{(r+a)(r+b)} dr. \end{aligned}$$

Substitute this formula into equation (1).

$$\begin{aligned} (1 - e^{2i\pi/3}) \int_\rho^R \frac{r^{1/3}}{(r+a)(r+b)} dr + \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz + \int_{C_R} \frac{z^{1/3}}{(z+a)(z+b)} dz \\ = 2\pi i \left[\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} + \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} \right] \end{aligned}$$

Take the limit now as $\rho \rightarrow 0$ and $R \rightarrow \infty$. The integral over C_ρ tends to zero, and the integral over C_R tends to zero. Proof for these statements will be given at the end.

$$(1 - e^{2i\pi/3}) \int_0^\infty \frac{r^{1/3}}{(r+a)(r+b)} dr = 2\pi i \left[\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} + \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} \right]$$

The multiplicities of $z + a$ and $z + b$ in the denominator are both 1, so the residues at $z = -a$ and $z = -b$ can be calculated by

$$\begin{aligned}\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} &= \phi_1(-a) \\ \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} &= \phi_2(-b),\end{aligned}$$

where $\phi_1(z)$ and $\phi_2(z)$ are the same function as $f(z)$ without the factors, $z + a$ and $z + b$, respectively.

$$\begin{aligned}\phi_1(z) = \frac{z^{1/3}}{z+b} &\Rightarrow \phi_1(-a) = \frac{(-a)^{1/3}}{-a+b} = \frac{(ae^{i\pi})^{1/3}}{-a+b} = -\frac{a^{1/3}e^{i\pi/3}}{a-b} \\ \phi_2(z) = \frac{z^{1/3}}{z+a} &\Rightarrow \phi_2(-b) = \frac{(-b)^{1/3}}{-b+a} = \frac{(be^{i\pi})^{1/3}}{-b+a} = \frac{b^{1/3}e^{i\pi/3}}{a-b}\end{aligned}$$

So then

$$\begin{aligned}\operatorname{Res}_{z=-a} \frac{z^{1/3}}{(z+a)(z+b)} &= -\frac{a^{1/3}e^{i\pi/3}}{a-b} \\ \operatorname{Res}_{z=-b} \frac{z^{1/3}}{(z+a)(z+b)} &= \frac{b^{1/3}e^{i\pi/3}}{a-b}\end{aligned}$$

and

$$\begin{aligned}(1 - e^{2i\pi/3}) \int_0^\infty \frac{r^{1/3}}{(r+a)(r+b)} dr &= 2\pi i \left[-\frac{a^{1/3}e^{i\pi/3}}{a-b} + \frac{b^{1/3}e^{i\pi/3}}{a-b} \right] \\ &= \frac{2\pi i}{a-b} e^{i\pi/3} (-a^{1/3} + b^{1/3}).\end{aligned}$$

Divide both sides by $1 - e^{2i\pi/3}$.

$$\begin{aligned}\int_0^\infty \frac{r^{1/3}}{(r+a)(r+b)} dr &= \frac{2\pi i}{a-b} \cdot \frac{e^{i\pi/3}}{1 - e^{2i\pi/3}} (-a^{1/3} + b^{1/3}) \\ &= \frac{2\pi i}{a-b} \cdot \frac{1}{e^{-i\pi/3} - e^{i\pi/3}} (-a^{1/3} + b^{1/3}) \\ &= \frac{2\pi i}{a-b} \cdot \frac{1}{[-2i \sin(\pi/3)]} (-a^{1/3} + b^{1/3}) \\ &= \frac{2\pi}{a-b} \cdot \frac{1}{\sqrt{3}} (a^{1/3} - b^{1/3})\end{aligned}$$

Therefore, changing the dummy integration variable to x ,

$$\boxed{\int_0^\infty \frac{\sqrt[3]{x}}{(x+a)(x+b)} dx = \frac{2\pi}{\sqrt{3}} \cdot \frac{\sqrt[3]{a} - \sqrt[3]{b}}{a-b} .}$$

The Integral Over C_ρ

Our aim here is to show that the integral over C_ρ tends to zero in the limit as $\rho \rightarrow 0$. The parameterization of the small circular arc in Figure 1 is $z = \rho e^{i\theta}$, where θ goes from 2π to 0 .

$$\begin{aligned} \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz &= \int_{2\pi}^0 \frac{(\rho e^{i\theta})^{1/3}}{(\rho e^{i\theta} + a)(\rho e^{i\theta} + b)} (\rho i e^{i\theta} d\theta) \\ &= \int_{2\pi}^0 \frac{\rho^{4/3}}{(\rho e^{i\theta} + a)(\rho e^{i\theta} + b)} (i e^{4i\theta/3} d\theta) \end{aligned}$$

Take the limit of both sides as $\rho \rightarrow 0$.

$$\lim_{\rho \rightarrow 0} \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz = \lim_{\rho \rightarrow 0} \int_{2\pi}^0 \frac{\rho^{4/3}}{(\rho e^{i\theta} + a)(\rho e^{i\theta} + b)} (i e^{4i\theta/3} d\theta)$$

The limits of integration are constant, so the limit may be brought inside the integral.

$$\lim_{\rho \rightarrow 0} \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz = \int_{2\pi}^0 \lim_{\rho \rightarrow 0} \frac{\rho^{4/3}}{(\rho e^{i\theta} + a)(\rho e^{i\theta} + b)} (i e^{4i\theta/3} d\theta)$$

Because of $\rho^{4/3}$ in the numerator, the limit is zero. Therefore,

$$\lim_{\rho \rightarrow 0} \int_{C_\rho} \frac{z^{1/3}}{(z+a)(z+b)} dz = 0.$$

The Integral Over C_R

Our aim here is to show that the integral over C_R tends to zero in the limit as $R \rightarrow \infty$. The parameterization of the large circular arc in Figure 1 is $z = R e^{i\theta}$, where θ goes from 0 to 2π .

$$\begin{aligned} \int_{C_R} \frac{z^{1/3}}{(z+a)(z+b)} dz &= \int_0^{2\pi} \frac{(R e^{i\theta})^{1/3}}{(R e^{i\theta} + a)(R e^{i\theta} + b)} (R i e^{i\theta} d\theta) \\ &= \int_0^{2\pi} \frac{R^{4/3}}{(R e^{i\theta} + a)(R e^{i\theta} + b)} (i e^{4i\theta/3} d\theta) \\ &= \int_0^{2\pi} \frac{R^{4/3}}{R^2 (e^{i\theta} + \frac{a}{R}) (e^{i\theta} + \frac{b}{R})} (i e^{4i\theta/3} d\theta) \\ &= \int_0^{2\pi} \frac{1}{R^{2/3}} \frac{1}{(e^{i\theta} + \frac{a}{R}) (e^{i\theta} + \frac{b}{R})} (i e^{4i\theta/3} d\theta) \end{aligned}$$

Take the limit of both sides as $R \rightarrow \infty$. Since the limits of integration are constant, the limit may be brought inside the integral.

$$\lim_{R \rightarrow \infty} \int_{C_R} \frac{z^{1/3}}{(z+a)(z+b)} dz = \int_0^{2\pi} \lim_{R \rightarrow \infty} \frac{1}{R^{2/3}} \frac{1}{(e^{i\theta} + \frac{a}{R}) (e^{i\theta} + \frac{b}{R})} (i e^{4i\theta/3} d\theta)$$

Because of $R^{2/3}$ in the denominator, the limit is zero. Therefore,

$$\lim_{R \rightarrow \infty} \int_{C_R} \frac{z^{1/3}}{(z+a)(z+b)} dz = 0.$$