

Problem 6

In each of Problems 1 through 12:

- Draw a direction field for the given differential equation.
- Based on an inspection of the direction field, describe how solutions behave for large t .
- Find the general solution of the given differential equation, and use it to determine how solutions behave as $t \rightarrow \infty$.

$$ty' + 2y = \sin t, \quad t > 0$$

Solution

The direction field is a two-dimensional vector field that shows what the direction of the solution is at every point in a region. Every solution to the differential equation is a curve drawn such that the direction field vectors are tangent to it at every point.

$$\langle dt, dy \rangle = \left\langle 1, \frac{dy}{dt} \right\rangle dt = \left\langle 1, \frac{-2y + \sin t}{t} \right\rangle dt$$




Figure 1: In blue are the direction field vectors and in red are possible solutions to the differential equation, depending what the initial condition is. All solutions appear to converge to $y = 0$ as $t \rightarrow \infty$.

$$ty' + 2y = \sin t$$

Divide both sides by t so that the coefficient of y' is 1.

$$y' + \frac{2}{t}y = \frac{1}{t}\sin t$$

This is a first-order linear inhomogeneous equation, so it can be solved with an integrating factor.

$$I(t) = \exp\left(\int^t \frac{2}{s} ds\right) = e^{2\ln t} = e^{\ln t^2} = t^2$$

Multiply both sides of the ODE by $I(t)$.

$$t^2y' + 2ty = t\sin t$$

The left side can be written as $d/dt(Iy)$ by the product rule.

$$\frac{d}{dt}(t^2y) = t\sin t$$

Integrate both sides with respect to t .

$$\begin{aligned} t^2y &= \int^t s \sin s ds + C \\ &= \sin t - t \cos t + C \end{aligned}$$

Divide both sides by t^2 to solve for $y(t)$.

$$y(t) = \frac{1}{t^2}(\sin t - t \cos t + C)$$

$y(t)$ vanishes as $t \rightarrow \infty$.

$$\lim_{t \rightarrow \infty} y(t) = 0$$