

Problem 9

In each of Problems 1 through 12:

- Draw a direction field for the given differential equation.
- Based on an inspection of the direction field, describe how solutions behave for large t .
- Find the general solution of the given differential equation, and use it to determine how solutions behave as $t \rightarrow \infty$.

$$2y' + y = 3t$$

Solution

The direction field is a two-dimensional vector field that shows what the direction of the solution is at every point in a region. Every solution to the differential equation is a curve drawn such that the direction field vectors are tangent to it at every point.

$$\langle dt, dy \rangle = \left\langle 1, \frac{dy}{dt} \right\rangle dt = \left\langle 1, \frac{-y + 3t}{2} \right\rangle dt$$




Figure 1: In blue are the direction field vectors and in red are possible solutions to the differential equation, depending what the initial condition is. In green is the asymptotic solution $y = 3(t - 2)$ that all other solutions tend to as $t \rightarrow \infty$.

$$2y' + y = 3t$$

Divide both sides by 2 to make the coefficient of y' 1.

$$y' + \frac{y}{2} = \frac{3}{2}t$$

This is a first-order linear inhomogeneous equation, so it can be solved with an integrating factor.

$$I(t) = \exp\left(\int^t \frac{1}{2} ds\right) = e^{t/2}$$

Multiply both sides of the ODE by $I(t)$.

$$e^{t/2}y' + \frac{1}{2}e^{t/2}y = \frac{3}{2}te^{t/2}$$

The left side can be written as $d/dt(Iy)$ by the product rule.

$$\frac{d}{dt}(e^{t/2}y) = \frac{3}{2}te^{t/2}$$

Integrate both sides with respect to t .

$$\begin{aligned} e^{t/2}y &= \int^t \frac{3}{2}se^{s/2} ds + C \\ &= 3e^{t/2}(t - 2) + C \end{aligned}$$

Divide both sides by $e^{t/2}$ to solve for $y(t)$.

$$y(t) = 3(t - 2) + Ce^{-t/2}$$

$y(t)$ tends to $3(t - 2)$ asymptotically as $t \rightarrow \infty$.