

Problem 13

In each of Problems 9 through 20:

- Find the solution of the given initial value problem in explicit form.
- Plot the graph of the solution.
- Determine (at least approximately) the interval in which the solution is defined.

$$y' = 2x/(y + x^2y), \quad y(0) = -2$$

Solution

Part (a)

This ODE is separable because it is of the form $y' = f(x)g(y)$, so it can be solved by separating variables.

$$\frac{dy}{dx} = \frac{2x}{y(1+x^2)}$$

Bring the terms with y to the left and bring the terms with x to the right.

$$y \, dy = \frac{2x}{1+x^2} \, dx$$

Integrate both sides.

$$\int y \, dy = \int \frac{2x}{1+x^2} \, dx \tag{1}$$

Make the following substitution in the integral in dx .

$$\begin{aligned} u &= 1 + x^2 \\ du &= 2x \, dx \end{aligned}$$

Equation (1) becomes

$$\begin{aligned} \frac{y^2}{2} &= \int \frac{du}{u} \\ &= \ln|u| + C \\ &= \ln(1+x^2) + C. \end{aligned}$$

Now apply the initial condition to determine C .

$$\frac{(-2)^2}{2} = \ln(1) + C \quad \rightarrow \quad 2 = C$$

As a result,


$$\begin{aligned} \frac{y^2}{2} &= \ln(1+x^2) + 2 \\ y^2 &= 2\ln(1+x^2) + 4 \\ y(x) &= \pm\sqrt{2\ln(1+x^2) + 4}. \end{aligned}$$

We choose the minus sign so that the initial condition is satisfied. Therefore,

$$y(x) = -\sqrt{2\ln(1+x^2) + 4}.$$

Part (b)

Below is a plot of $y(x)$ versus x .

**Part (c)**

The solution we found is only valid along the curve that passes through $x = 0$ (the y -axis) because $x = 0$ is where the initial condition is given. It is valid over $-\infty < x < \infty$, the domain of y .