

## Problem 17

In each of Problems 9 through 20:

- Find the solution of the given initial value problem in explicit form.
- Plot the graph of the solution.
- Determine (at least approximately) the interval in which the solution is defined.

$$y' = (3x^2 - e^x)/(2y - 5), \quad y(0) = 1$$

---

### Solution

#### Part (a)

This ODE is separable because it is of the form  $y' = f(x)g(y)$ , so it can be solved by separating variables.

$$\frac{dy}{dx} = \frac{3x^2 - e^x}{2y - 5}$$

Bring the terms with  $y$  to the left and bring the terms with  $x$  to the right.

$$(2y - 5) dy = (3x^2 - e^x) dx$$

Integrate both sides.

$$\int (2y - 5) dy = \int (3x^2 - e^x) dx$$
$$y^2 - 5y = x^3 - e^x + C$$

Now apply the initial condition to determine  $C$ .

$$1^2 - 5(1) = 0 - 1 + C \quad \rightarrow \quad C = -3$$

As a result,

$$y^2 - 5y = x^3 - e^x - 3$$
$$y^2 - 5y - (x^3 - e^x - 3) = 0$$
$$y(x) = \frac{5 \pm \sqrt{25 + 4(x^3 - e^x - 3)}}{2}.$$

We choose the minus sign so that the initial condition is satisfied.


$$y(x) = \frac{5 - \sqrt{25 + 4(x^3 - e^x - 3)}}{2}$$

Therefore,

$$y(x) = \frac{5 - \sqrt{13 + 4(x^3 - e^x)}}{2}.$$

**Part (b)**

Below is a plot of  $y(x)$  versus  $x$ .

**Part (c)**

The solution we found is only valid along the curve that passes through  $x = 0$  (the  $y$ -axis) because  $x = 0$  is where the initial condition is given. It is valid over the domain of  $y$ .

$$25 + 4(x^3 - e^x - 3) \geq 0$$

Judging from the graph, it is  $-1.44 \lesssim x \lesssim 4.63$ .