

Problem 20

In each of Problems 17 through 22, find the solution of the given initial value problem. Sketch the graph of the solution and describe its behavior for increasing t .

$$y'' + y = 0, \quad y(\pi/3) = 2, \quad y'(\pi/3) = -4$$

Solution

Since this is a linear homogeneous constant-coefficient ODE, the solution is of the form $y = e^{rt}$.

$$y = e^{rt} \rightarrow y' = re^{rt} \rightarrow y'' = r^2e^{rt}$$

Substitute these expressions into the ODE.

$$r^2e^{rt} + e^{rt} = 0$$

Divide both sides by e^{rt} .

$$r^2 + 1 = 0$$

$$r = \{-i, i\}$$

Two solutions to the ODE are $y = e^{-it}$ and $y = e^{it}$, so the general solution is a linear combination of the two.

$$\begin{aligned} y(t) &= C_1e^{-it} + C_2e^{it} \\ &= C_1[\cos(-t) + i\sin(-t)] + C_2[\cos(t) + i\sin(t)] \\ &= C_1(\cos t - i\sin t) + C_2(\cos t + i\sin t) \\ &= C_1\cos t - iC_1\sin t + C_2\cos t + iC_2\sin t \\ &= (C_1 + C_2)\cos t + (-iC_1 + iC_2)\sin t \end{aligned}$$

Using C_3 for $C_1 + C_2$ and C_4 for $-iC_1 + iC_2$, the real general solution is

$$y(t) = C_3\cos t + C_4\sin t.$$

Take a derivative of it.


$$y'(t) = -C_3\sin t + C_4\cos t$$

Apply the initial conditions now to determine C_3 and C_4 .

$$\begin{aligned} y\left(\frac{\pi}{3}\right) &= C_3\left(\frac{1}{2}\right) + C_4\left(\frac{\sqrt{3}}{2}\right) = 2 \\ y'\left(\frac{\pi}{3}\right) &= -C_3\left(\frac{\sqrt{3}}{2}\right) + C_4\left(\frac{1}{2}\right) = -4 \end{aligned}$$

Solving this system of equations yields $C_3 = 1 + 2\sqrt{3}$ and $C_4 = -2 + \sqrt{3}$. Therefore,

$$y(t) = (1 + 2\sqrt{3})\cos t + (-2 + \sqrt{3})\sin t.$$



The solution oscillates with a constant amplitude and period.