

## Problem 1.11

The position of a moving particle is given as a function of time  $t$  to be

$$\mathbf{r}(t) = \hat{\mathbf{x}}b \cos(\omega t) + \hat{\mathbf{y}}c \sin(\omega t),$$

where  $b$ ,  $c$ , and  $\omega$  are constants. Describe the particle's orbit.

---

### Solution

Write the given position vector as

$$\begin{aligned}\mathbf{r}(t) &= \langle b \cos \omega t, c \sin \omega t \rangle \\ &= \langle x(t), y(t) \rangle.\end{aligned}$$

To determine the particle's orbit, it's necessary to find the equation satisfied by  $x$  and  $y$ . Start with a known trigonometric identity involving  $\cos \omega t$  and  $\sin \omega t$ .

$$\begin{aligned}\cos^2 \omega t + \sin^2 \omega t &= 1 \\ \frac{b^2 \cos^2 \omega t}{b^2} + \frac{c^2 \sin^2 \omega t}{c^2} &= 1 \\ \frac{x^2}{b^2} + \frac{y^2}{c^2} &= 1\end{aligned}$$

This is the equation for an ellipse, so the particle's orbit is elliptical. At  $\omega t = 0$ , the position vector is

$$\mathbf{r}(0) = \langle b, 0 \rangle$$

so the particle starts on the  $x$ -axis at  $(b, 0)$  and moves around the ellipse at angular speed  $\omega$ .