

Exercise 13

Use the *Laplace transform method* to solve the Volterra integral equations:

$$u(x) = 2e^x - 2 - x + \int_0^x (x-t)u(t) dt$$

Solution

The Laplace transform of a function $f(x)$ is defined as

$$\mathcal{L}\{f(x)\} = F(s) = \int_0^{\infty} e^{-sx} f(x) dx.$$

According to the convolution theorem, the product of two Laplace transforms can be expressed as a transformed convolution integral.

$$F(s)G(s) = \mathcal{L}\left\{\int_0^x f(x-t)g(t) dt\right\}$$

Take the Laplace transform of both sides of the integral equation.

$$\begin{aligned} \mathcal{L}\{u(x)\} &= \mathcal{L}\left\{2e^x - 2 - x + \int_0^x (x-t)u(t) dt\right\} \\ U(s) &= 2\mathcal{L}\{e^x\} - 2\mathcal{L}\{1\} - \mathcal{L}\{x\} + \mathcal{L}\left\{\int_0^x (x-t)u(t) dt\right\} \\ &= 2\mathcal{L}\{e^x\} - 2\mathcal{L}\{1\} - \mathcal{L}\{x\} + \mathcal{L}\{x\}U(s) \\ &= 2\left(\frac{1}{s-1}\right) - 2\left(\frac{1}{s}\right) - \frac{1}{s^2} + \left(\frac{1}{s^2}\right)U(s) \end{aligned}$$

Solve for $U(s)$.

$$\begin{aligned} \left(1 - \frac{1}{s^2}\right)U(s) &= \frac{2}{s-1} - \frac{2}{s} - \frac{1}{s^2} \\ (s^2 - 1)U(s) &= \frac{2s^2}{s-1} - 2s - 1 \\ (s+1)(s-1)U(s) &= \frac{s+1}{s-1} \\ U(s) &= \frac{1}{(s-1)^2} \end{aligned}$$

Take the inverse Laplace transform of $U(s)$ to get the desired solution.

$$\begin{aligned} u(x) &= \mathcal{L}^{-1}\{U(s)\} \\ &= \mathcal{L}^{-1}\left\{\frac{1}{(s-1)^2}\right\} \\ &= xe^x \end{aligned}$$